

3-Thiocyanato- 1H- indoles as potential anticancer agents: Two dimensional quantitative structure activity relationship study

Mr.Sivakumar Peta ,Dr.M.Soujanya , Mrs.S.Usharani , Mr.B.Kondalrao

Department of Pharmaceutical Chemistry,

Email ID: sivapeta369@gmail.com , Mobile no: 9494144800

GOKULA KRISHNA COLLEGE OF PHARMACY

Abstract


We conducted two-dimensional quantitative structure activity relationship (2D QSAR) research on a new series of 3-thiocyanato-1H-indoles in an effort to identify powerful anti-cancer drugs. variety of 3-thiocyanato-1H-indoles were subjected to 2D-QSAR using Vlife MDS 4.3. The k-nearest neighbors (kNN) approach, used to Vlife molecular design suites (MDS), yielded a statistically verified two-dimensional quantitative structure activity relationship model. Cytotoxicity activity against the HL60 human cancer cell line was associated with Model 3 statistical data ($q^2 = 0.8001$, $\text{pred } r^2 = 0.4082$). The LOO approach was used for validation. Final Thoughts: The model now includes three attributes that positively correlate with the cytotoxicity activity. There is hope that novel, more effective anticancer drugs could be developed using this proven 2D QSAR model.

Keywords: 2-dimensional quantum search for anticancer drugs using regression analysis; 3-thiocyanato-1H-indoles; HL60 cell line.

Introduction

The unique capacity of the compounds produced by heterocyclic chemistry to bind reversibly to proteins and imitate the structure of peptides makes it a very useful source of new molecules with various biological functions. (1) to four (3) Indole, also known as benzopyrrole, is a heterocyclic compound with one nitrogen atom (N) substituted for one carbon atom in the ring. As a privileged structure that binds to several receptors with high affinity, the indole moiety is widespread and ranks among the most prevalent heterocycles among physiologically active natural compounds, medicines, and agrochemicals (5). The therapeutic implications of Indole have been highlighted in published publications as follows: anti-viral, anti-depressant, anti-hyperlipidemic, anti-inflammatory, anti-psychotic, anti-microbial, anti-oxidants, anti-HIV, immunomodulator, anti-leukemia, (19).(21-22) Natural substances with strong pharmacodynamic Indole nucleus activity include reserpine, bufotenine, tryptophan, serotonin, vinblastine, vincristine, tryptamine derivatives, and others.

As the second-biggest killer of humans, cancer poses a serious danger to human health. (29-32) The World Health Organization (WHO) projects that 12 million people will lose their lives to cancer by the year 2030. (33) radiation and chemotherapy are two of the current cancer therapies, however the most remarkable pharmaceutical approach to cancer would still be a combination of radiation and significant surgery. The limitations of the current anticancer drugs, including as their toxicity to normal cells and acquired tumor resistance, persist despite ongoing research. Thus, enhancing the pharmacological profile and conducting cutting-edge cancer research depend on the discovery of effective, safe, and selective anticancer agents. (34) For the purpose of predicting biological activities, especially in medication design, the quantitative structure-activity relationship (QSAR) method became very helpful and extensively used. The premise upon which this method rests is that alterations to their biological activity may be associated with changes to their chemical structures. (35 and 36) Margiani et al. prepared a battery of 3-thiocyanato-1H-indoles and tested their cytotoxic effects on several cancer cell lines. Anticancer activities with improved treatment safety and effectiveness were the goals of this work, which intended to clarify the structural properties of 3-thiocyanato-1H-indole derivatives. (37)




Material and Methods

Computer aided drug designing based on quantitative structure activity relationship (QSAR) is the study of the quantitative relationship between the experimental activity of a set of compounds and their physicochemical properties. Different statistical approaches are used while building a QSAR model, where experimental information associated with biological activity, is used as dependent variable.⁽³⁵⁻³⁶⁾ In the present study, all computational work (2D- QSAR) was performed using Vlife MDS 4.3 QSAR plus software on a HP Pentium IV 2.80 GHz Processor / Microsoft Win XP Home Edition system.

2D-QSAR modelling and dataset: Information on cytotoxicity-inducing activities Data from published sources were used to determine the IC₅₀ (μM) of the twenty compounds listed in Table 1.⁽³⁷⁾ With the use of the Mosamann technique in an MTT test, the experimental IC₅₀ values were determined for HL-60 cell lines; IC₅₀ represents a 50% suppression of cancer development. Chemicals were deemed active if their pIC₅₀ values were between 5.25 and 6.20. Table 2 lists the dependent variable for 2D investigations, which is the negative logarithm of the observed IC₅₀ (μM) [pIC₅₀ = -log (IC₅₀)]. Chemdraw Ultra 8.0 was used to build 2D structures, which were then transformed into 3D ones using the same program. Molecular mechanics (MMFF) and Montocarlo conformational search were used to minimize energy and analyze conformations of all three-dimensional molecules until the RMS gradient value was less than 0.001 kcal/mol Å. In order to determine the physicochemical properties of stable molecules, this Montocarlo conformational search approach produces a number of different conformations after energy minimization. After initially calculating 240 2D descriptors using energy-minimizing geometry, the number was subsequently decreased to 156 using invariable column selection. The following steric, topological, electronic, molecular, and structural descriptors were computed: chiV6chain, chiV4pathcluster, Idw Average (steric), chi1, chi5, chiV0, chiV3, chiV4, chiV5, 3PathCount, chi6chain, chiV6chain, chiV3Cluster, 3ClusterCount, chi4pathCluster, chiV4pathCluster, 4pathClusterCount, kappa3, k1alpha, k2alpha, VChi 3 cluster, VChi 4 cluster, and VChi 5 path. The Kier symmetry index, the alpha 1 and alpha 3 shapes, and the Kier shape 2 Ch0, Ch2, and Chi 3

group, not Here are some terms related to Chi-square: VChi-1, VChi-3, VChi-4, VChi-4 path/cluster, Kier-1, Kier-3, Kier-alpha-2, Kier-flex, and Kier steric descriptor. Charge index 1, charge index 3, charge index 5, charge index 7, charge index 9, valence charge index 2, valence charge index 4, valence charge index 6, valence charge index 8, valence charge index 10, delta chi 0, delta chi 2, delta chi 3 cluster, delta chi 5 path, variation chi 1, variation chi 3, variation chi 5, delta chi 1, delta chi 3 path, delta chi 4 path/cluster, variation chi 0, variation chi 2, bound charge index 2, k2alpha With a topological chi-five route and zero path count, Various terms such as maximum positive charge, maximum positive hydrogen charge, relative negative charge, hydrophobic SA-MPEOE, negative charged polar, maximum negative charge, charge polarization, polarity parameter, and relative positive charge are used. surface area with a positive potential, surface area with a negative potential, average potential, average +ve potential, average -ve potential, distance between the most positive and negative potential, average -ve potential (electronic), and so on.

Table 1: Series of 3-thiocyanato-1H-indole derivatives



Compound no.	R1	R2	R3
1	H	H	H
2	H	H	4-Me-C ₆ H ₄
3	H	H	OMe
4	H	H	Br
5	H	H	CN
6	Ph	H	H
7	Me	H	4-Me-C ₆ H ₄
8	4-MeO-C ₆ H ₄	H	H
9	Me	H	H
10	4-Cl-C ₆ H ₄	H	H
11	H	C ₆ H ₅	H
12	H	4-Me-C ₆ H ₄	H
13	H	4-MeO-C ₆ H ₄	H
14	H	3CF ₃ -C ₆ H ₄	H
15	H	4-Cl-C ₆ H ₄	H
16	Me	C ₆ H ₅	H
17	Me	4-Me-C ₆ H ₄	H
18	Me	4-MeO-C ₆ H ₄	H
19	Me	3CF ₃ -C ₆ H ₄	H
20	Me	4-Cl-C ₆ H ₄	H

Table 2: Observed and Predicted activities of compounds

Compound no.	Actual/Observed Activity	Predicted Activity
1	5.49	5.38
2	6.59	6.17
3	5.25	5.51
4	5.52	5.37
5	5.25	5.45
6	5.84	5.92
7	5.39	5.52
8	5.85	5.85
9	5.81	5.85
10	5.45	5.49
11	6.20	6.06
12	5.64	6.18
13	5.35	6.08
14	6.16	6.18
15	5.97	5.84
16	5.86	5.86
17	5.82	5.89
19	5.59	5.42
20	5.96	5.85

Selection of training and test set: A data set of 20 molecules belonging to 3-thiocyanato-1H-indoles derivatives as

anticancer agents were taken from the literature⁽³⁷⁾ and used for QSAR study. The dataset was divided into training set (12 compounds) and test set (8 compounds) by Sphere Exclusion (SE) method for principal component regression (PCR), multiple linear

regression (MLR), partial least squares (PLS), k-nearest neighbours (kNN) model. In classical sphere exclusion algorithm, each selected molecule generates a hyper- sphere around itself, where any molecule inside the sphere is excluded from the selection in the train set and driven towards the test set. The number of compounds selected and the diversity among them can be determined by adjusting the radius of the sphere (R). Inhibitory activity i.e. pIC₅₀ has been considered as dependent variable and the remaining descriptors as independent parameters.

Regression analysis: Quantitative Structure-Activity Relationship (QSAR) analysis is a parametric method that researchers use to learn about the action mechanism and key structural components of a class of drugs. Specifically, 20 compounds were tested for their biological activity using regression analysis, PCR, MLR, PLS, and kNN as statistical methods. Correlative parameters were chosen based on their physicochemical or structural properties. Using a cross-correlation threshold of 0.7, four variables were included in the final equation for PCR, MLR, and PLS. The term selection criteria included r², F-test 'in,' at 4, and 'out,' at 3.99, in addition to r² and the F-test. We set the scaling to auto scaling, set the variance cut off to 0.1, and set the number of random iterations to 10. For the purpose of evaluating QSAR models, the following statistical measures were utilized: n, the number of chemicals in regression, r, the number of descriptors in a model, k, and the F-test (Fisher test value) for statistical significance. Equations (F), (q²), (pred_r²), (r² se) and (q² se) represent the standard error (SE) of estimate, as well as the cross-validated correlation coefficient and the predictive squared correlation coefficients. The relative fit of the regression equation is measured by the regression coefficient r². For a better regression fit, the values of the correlation coefficient should be closer to 1.0, the significance level. The ratio of the variation explained by the model to the variance owing to the error in the regression is shown in the F-test. A statistically significant model is one with a high F-test score. One of the validation parameters, predictive r² (r²_pred), was computed to assess the model's predictive ability; a value higher than 0.5 indicates that the QSAR model has strong predictive capacity. The biological activity was associated with the physico-chemical descriptor values using a variety of statistical models that were created using PCR, MLR, PLS, and kNN based regression techniques combined with forward, forward backward, genetic algorithm, and simulated annealing method.

Validation of QSAR model: Model validation is done to analyse the internal stability and predictive ability of the QSAR models. The best to evaluate quality of regression model is internal and external validation. Internal validation is carried out using leave one out (q²,

LOO) method. For calculating q² each molecule in the training set was eliminated once from training set and the activity of eliminated molecule was predicted by using model developed by the remaining molecules. This q² described the internal stability of the model⁽³⁹⁾ and is expressed as shown in equation (Eq A.1)

$$q^2 = 1 - \frac{\sum (Y_{\text{pred}} - Y_{\text{act}})^2}{\sum (Y_{\text{act}} - Y_{\text{mean}})^2} \quad (\text{Eq A.1})$$

Secondly, the predictive ability of the model i.e. external validation was confirmed by comparing the observed value of the test set molecules with predictive value of test molecules and is indicated by predicted r² as shown in equation (Eq A.2).

$$\text{pred}_r^2 = 1 - \frac{\sum (Y_{\text{pred}}(\text{Test}) - Y_{\text{Test}})^2}{\sum (Y_{\text{Test}} - Y_{\text{Training}})^2} \quad (\text{Eq A.2})$$

Results

Selected molecular parameters computed for all the 20 molecules were used to develop QSAR equation by relating their corresponding inhibitory activities i.e. IC₅₀ values. Aim was to establish a predictive model with a number of logical descriptors to get good generalization performance and which can be further utilized for the synthesis of 3-thiocyanato-1H-indoles as anticancer agents. Various models were developed using, PCR, MLR, PLS and kNN and the model having best fit with minimum number of descriptors has considered or found to be the best model. When this point is achieved, no further considerable improvement in the regression coefficient (r² and q²) values were observed even if a new descriptor is added. In the present study PCR, MLR, PLS and kNN, methods employing three, four or more variable combinations for combined dataset generated around 300, equations out of which, the reasonable acceptable ones were selected for discussion. The different models generated for better correlation by statistical analysis have been given in **Table 3**, whereas **Table 4** contains various descriptors, their categories as well as their contribution towards each descriptor.

Table 3: Various significant models obtained by MLR and kNN methods and their statistical parameters

Model	Equation	N	r ²	q ²	Pred r ²	F-test
MLR (SA)	Eq B.1	14	0.9393	0.6493	0.3052	4.6419
kNN (SA)	Eq C.1	15	-	0.7224	0.3867	-
kNN (SA)	Eq D.1	16	-	0.8001	0.4082	-

Table 4: Various descriptors obtained by MLR method (Eq B.1); their categories, contributions and significance

Descriptor	Category	Contribution	Significance
Average -ve Potential	Electrostatic	+	total -ve electrostatic potential on van der Waals surface area of the molecule
Bromine Count	Element count	+	number of bromine atoms in a compound
SsCH3Count	Estate Numbers	-	total number of -CH ₃ group connected with single bond
Idw Average	Information Theory Based	+	information-based descriptors
SA Hydrophilic Area	Hydrophobicity	-	vd W surface descriptor showing hydrophobic surface area
-Ve Potential Surface Area	Electrostatic	-	total van der Waals surface area with negative electrostatic potential of the molecule
SK Most Hydrphobic Hydrophilic Distance	Hydrophobicity	-	total van der Waals surface area with negative electrostatic potential of the molecule
SA Most Hydrphobic Hydrophilic Distance	Hydrophobicity	+	distance between most hydrophobic and hydrophilic point on the vdW surface
XK Most Hydrophobic Hydrophilic Distance	Hydrophobicity	+	distance between most hydrophobic and hydrophilic point on the vdW surface
+ve Potential Surface Area	Electrostatic	+	total van der Waals surface area with positive electrostatic potential of the molecule

Eq B.1 represents the model 1, obtained by MLR regression analysis

$pIC_{50} = +13.9818$ (Average-ve Potential) $+0.0709$ (Bromine Count) -0.1108 (SsCH₃Count) $+1.5159$ (Idw Average) -0.0574 (SA Hydrophilic Area) -0.0163 (-ve Potential Surface Area) -0.0277 (SK Most Hydrphobic Hydrophilic Distance) $+0.0842$ (SA Most Hydrphobic Hydrophilic Distance) $+0.1995$ (XK Most Hydrophobic Hydrophilic Distance) $+0.0200$ (+ve Potential Surface Area) **(Eq B.1)**

Contributions of the various descriptors, fitness plots and actual and predicted of the training and sets molecules w.r.t. MLR model are shown in **Fig. 1, 2**. Model with low pred r^2 0.3052 (31%), and cross- validated correlation coefficient q^2 0.6493 was not found to be satisfactory for a good correlation between the structure and activity.

k-Nearest Neighbour method using simulated annealing (SA) was performed in order to get improved QSAR models, kNN method is used for modelling linear relationship between a dependent variable Y (pEC₅₀) and independent variable X (2D descriptors). In this method, an unknown pattern is classified according to the majority of the class membership of its k nearest neighbors in the training set. The nearness is measured by an appropriate distance metric (e.g. a molecular similarity measure, calculated using field interactions of molecular structures).⁽⁴⁰⁾ Using same data set, a new kNN based model 2 (Eq C.1) was generated.

$$pIC_{50} = 0.0809 \text{ 0.0845}(\text{SaasN}(\text{N oxide})\text{count}) + 1.0000$$

$$1.0000(\text{XK Hydrophobic Area}) + 362.3820 \text{ 366.7580}(\text{SssO count}) \quad (\text{Eq C.1})$$

Generated QSAR model 2 (kNN method) shows high cross-validation correlation coefficient, q^2 between descriptors (SaasN(N oxide)count, XK Hydrophobic Area, SssO count) and cytotoxicity

activity, but the low r^2 value 0.3867 i.e. predicted correlation coefficient, indicates the model is not significant. **Fig. 3** shows the fitness plot and actual & predicted activity of the training and test molecules.

The compound 18 was found to be an outlier and its omission resulted into model 3 with improved correlation i.e. **Eq. D.1** and was found to be the best model that exhibits good external pre-dictivity indicated by pred r^2 0.4082 and cross-validated correlation coefficient q^2 0.8001 using kNN by simulated annealing method. The fitness plot and of observed, predicted activities of the 20 molecules have been given in **Fig. 4**.

$$pIC_{50} = 0.0000 \text{ 0.0000}(\text{H-Donor Count}) + 360.7880$$

$$361.3780(\text{SA Hydrophobic Area}) + 9.2460 \text{ 9.2570}(\text{StNE-index}) \quad (\text{Eq D.1})$$

The best model is indicating the effect of H-Donor Count, SA Hydrophobic Area and StNE-index on the biological activity. H-Donor Count an individual descriptor defines the number of hydrogen bond donor, but in the present studies no significant correlation has been found with biological activity as indicating in the equation with its value ranging from 0.0000 0.0000. SA Hydrophobic Area descriptor is a hydrophobic Slog P descriptor describing the hydrophobic surface area, and present equation has indicated its positive correlation (360.7880 361.3780) with the biological activity. It means higher the hydrophobic surface area higher would be the anticancer activity. The importance of number of nitrogen atom connected with one triple bond is further indicated by StNE-index descriptor, an estate contribution descriptor. Descriptor has been found to be positive correlated with the biological activity with value ranging from 9.2460 - 9.2570 thus greater contribution of StNE-index results into greater cytotoxicity against cancer cell line HL60.

Table 5: Various descriptors obtained by kNN (Eq C.1) method; their categories, contributions and significance

Descriptor	Category	Contribution	Significance
SaasN(N oxide)count	Estate Numbers	+	total number of nitro oxide group connected with one single along with two aromatic bonds
K Hydrophobic Area	Hydrophobicity X log pK	+	total number of oxygen connected with two single bonds
SssO count	Estate Number	+	total number of oxygen connected with two single bonds

Table 6: Various descriptors obtained by kNN (Eq D.1) method; their categories, contributions and significance

Descriptor	Category	Contribution	Significance
H-Donor Count	Individual	+	number of hydrogen bond donor atoms
SA Hydrophobic Area	Hydrophobicity Slog pA	+	descriptor showing hydrophobic surface area

StNE-index	Estate contributions	+	number of nitrogen atom connected with one triple bonds
------------	----------------------	---	---------------------------------------------------------

Conclusions

To test their potential anticancer effects, 20 compounds of 3-thiocyanato-1H-indoles were run through a 2D-QSAR. Results from the current series of 3-thiocyanato-1H-indoles shown high correlation, as shown by the best model 3 with $q^2 = 0.8001$ and $\text{pred } r^2 = 0.4082$ as anticancer activity against cell line (HL-60). All of the suggested QSAR models were statistically significant. The QSAR model's predictive capabilities were examined using the LOO approach, which yields impressive internal predictivity. Numerous physicochemical characteristics were shown to have an effect on the cell line (HL-60) in the QSAR investigation. These included the SA Hydrophobic Area (which had a positive contribution), the StNE-index (which had a positive contribution), and H-Donor Count (which had no effect). These findings will be useful in the development of novel anticancer drugs derived from 3-thiocyanato-1H-indoles.

Conflict of interest

The authors confirm that this article content has no conflict of interest.

Acknowledgment

The authors are thankful to the Dr Amit Bedi, and Dr Kundan; Vlife Science Technologies Pvt. Ltd, Pune, India, for providing us with the Vlife MDS 4.3 software and for their continual support in the QSAR study. The grants received from Department of Science and Technology (DST), are gratefully acknowledged.

References

1. "Horton DA, Bourne GT, and Smythe ML." 2003. Published. The synthesis via combinatorial means of privileged substructures or bicyclic privileged structures. *Chemistry Review* 103:893-930.
2. The authors of the 2011 study were Regina LG, Coluccia A, Brancale A, Piscitelli F, Gatti V, Maga G, Samuele A, Pannecouque C, Schols D, Balzarini J, Novellino E, and Silvestri R. New cyclic substituents at indole-2-carboxamide allow indolylarylsulfones to block HIV-1 non-nucleoside reverse transcriptase. In the *Journal of Medical Chemistry*, volume 54, pages 1587–1598.
3. Roberts M, Cosset FL, Polyak SJ, Boriskin YS, Molle J, Pecheur EI, Lavillette D, Alcaras F, 2007. The biochemical process by which the broad-spectrum antiviral arbidol inhibits the hepatitis C virus. *Journal of Biochemistry*, 46, 6050–6059.
4. Musella et al. (2016) conducted the study. An indol-based derivative has been identified as a very effective and specific inhibitor of varicella zoster virus (VZV). *European Journal of Medical Chemistry*, 124:773–781.
5. In 2015, Saini, Kumar, and Narasimhan published a thesis. An Overview of Indole Derivatives' Activities in the Central Nervous System. *Heart Rhythm Drugs Citation: Chem.* 16:19-28.
6. The authors of the 2013 study were Shattat G, Qirim AT, Sheikha GA, Hiari AY, Sweidan K, Qirim AR, Hikmat S, Hamadneh L, and Kouz AS. Investigation into the impact on the plasma lipid profile of Triton-WR-1339-induced Wistar rats of new 5-fluoro-N-(9,10-dihydro-9,10-dioxoanthracen-8-yl)-1H-indole-2-carboxamide compounds was conducted. *Enzyme inhibitory medicinal chemistry* 28:863-869.
7. A Mini Review of Indole as a Core Anti-Inflammatory Agent—Hemalathaa K, Madhumitha G. Selvaraj MR. 2013. *Chemistry Research Letters*, 2:287-292.
8. In 2011, Galadino SL, Pitta IR, Goncalves ST, Maia MB, Colaco NC, Galdino AS, Malta DJ, Laranjeira LP, and Lima DMC were all involved. Indole-imidazolidine derivatives: anti-inflammatory and antinociceptive actions. The publication is *Int Immunopharmacol* and the page numbers are 11:1816–1822.
9. By 2014. 9. Alka B, Umesh S, and Tania P. Complete the synthesis, docking, and pharmacological testing of new atypical antipsychotics based on indole. Published in the *European Journal of Medical Chemistry*, volume 74, pages 477–490.
10. In 1999, Denschenes, Lin, Ault, and Fassler published a study. An analysis of three potential histidine kinase inhibitors for bacteria and their antifungal implications. *Pharmaceutical Agents and Chemotherapy*, 43, 1700–1703.
11. Voelter W., Hiari YMA, Qaisi AM, and Abadelah MM (2006). 11. Antimicrobial and synthetic studies on 3-(aryl)-and 3-(heteroaryl) indoles with various substituents. *Monthly Journal of Chemistry*. 137:243-248.
12. In 2010, Monica ES, Luisa CC, Ribeiro D, Couto D, Freitas M, Gomes A, Ferreira LM, Fernandes E, Manuel M, and Marques B. conducted a study on the antioxidant activity of synthesising and screening undiscovered indole derivatives. Article published in the *European Journal of Medical Chemistry*, volume 45, pages 4869–4878.
13. Morse GD, Reichman RC, Fischl MA, Para M, Leedom J, Powderly W, Demeter LM, Resnick L, Bassiakos Y, Timpone J, Cox S, Batts D. 2010. Ateviridine mesylate concentration-targeted phase I trials in HIV-infected patients: dosing recommendations and pharmacokinetic investigations. *Virus Research* 45:47–58.
14. Leadicer D. *Organic Synthesis of Drugs*, 7th Edition, pages 41–154.
15. The year 2009, Prince HM. T cell lymphoma-active new pan-deacetylase inhibitor, Bishton M. *Hematology Meeting Abstracts* 3:33–38.

16. Mack GS. (2010). 16. Beyond selection, there is more. *Natural Biotechnology*, 28, 1259–1266, 2017.
17. 17. Manju SL with Sravanthi TV in 2016. 8. An intriguing framework for the creation of drugs—indoles. Chilton WS, Bigwood J, Gensen RE. 1979. *Eur. J. Med. Chem.* 91:1–10. Historical and biosynthetic observations on psilocin, bufotenine, and serotonin. *Journal of Psychedelic Drugs*, 11(1), 61–69.
18. 19. 1995, Radwanski ER, Last RL. Biochemical and molecular genetics of tryptophan metabolism and production. *The journal Plant Cell*, volume 7, pages 921–934, 2016.
19. 9. Berger, M., Gray, JA, and Roth, BL. Understanding serotonin's broader biological role... *Medical History Review*, 60, 335–366.
20. The bioproduction and effects on human health of indole alkaloids from *Catharanthus roseus* were discussed in a 2015 study by Almagro, Fernandez, and Pedreno. *chemicals*. 20:2973-3000.
21. The indole alkaloid marine natural products are a promising new avenue for the treatment of cancer, neurological disorders, and parasites (Waseem G., Hamann MT., 2005). (*Life Sciences*, 78, 442-453).
22. 23. Bach PB, Jett JR, Pastorino U, Tockman MS, Swensen SJ, Begg CB. 2007. Screening with computed tomography and outcomes for lung cancer. *24-JAMA*, 297:953-961. Jones, J.B. (2000). Identifying targets and developing drugs for cancer research based on mechanisms. *Science*, 287, 1969–1973, 25 minutes. From 2006, Lars V., Tobias V., and Herbert W. *Biological and Chemical Spatial Navigation: Using PSSC and SCONP as Developing Compound Collections on Scaffolds of Natural Products*. 25: 449 in *QSAR Combin. Sci.* 26. Published in 2008 by Rojo F, Albanell J, Rovira A, Corominas JM, and Manzarbeitia F. Targeted therapy in breast cancer. *Scientific Diagnostics and Therapeutics*. 25:245-261. 27. For 2008, the authors were Ruiz FM, Gil RR, Morreale A, Ortiz AR, Fabrega C, and Bravo J. Novel nucleosidic DNA alkyltransferase inhibitors discovered using structure-based approaches: virtual screening, in vitro and in vivo experiments. *Journal of Chemical Inf. Modeling*, 48, 844–854. 28. Published in 2003 by Barrett CT and Barrett JF. Is the expanded range of antibacterials sufficient to address the growing issue of resistance? Report in *Biotechnology: Current Opinion*, 4:621-626, 29. In 2005, Richard AL. An Adaptable Approach to Utilizing QSAR Models for Lead Optimization. 30-*Journal of Medical Chemistry*, 48:1638-1648. With contributions from Andreas B. Zhan, John WD, Meir G., and James HN. *Integrating Chemical and Biological Domains: "Target Fishing" with Two- and Three-Dimensional Molecular Characteristics*
23. Citation: Margiani PF, Paulo BNS, Teresinha GS, Teodoro SK, Gardenia CGM, Claudio CS. 2016. *Journal of Medical Chemistry*, 49: 6802-6810. The development and first assessment of 3-thiocyanato-1H-indoles as possible anticancer medicines. Published in the *European Journal of Medical Chemistry*, volume 118, pages 21–26, 32. The authors of the 1990 study were Skehan, Storeng, Scudiero, Monks, McMahon, Vistica, Warren, Bodesch, Kenney, and Boyd. A novel colorimetric cytotoxicity test has been developed for the purpose of medication screening against cancer. *Journal of the National Cancer Institute*, 82: 1107–1112. b) Mossman, T. 1983. Fast colorimetric method for assessing cell viability and proliferation; use in cytotoxicity and proliferation tests. *Journal of Immunolog. Methods*. 65:55-63.