



Comprehensive Strategies and Emerging Trends in the Global Fight Against Malaria

Dr. S. Peer Mohamed

**Assistant professor, Department of Zoology, Sadakathullah Appa College (Autonomous),
Rahmath Nagar, Tirunelveli-627011, Affiliated to Manonmaniam Sundaranar University,
Tirunelveli, Tamil Nadu, India**

Abstract:

Malaria remains a significant global health challenge, particularly in regions where poverty, climate, and limited access to healthcare converge. This review article provides a comprehensive overview of current research findings, control strategies, and emerging trends in the fight against malaria. Drawing from a wide range of scientific literature and field reports, we delve into the intricate web of factors influencing malaria transmission, its impact on communities, and the innovative approaches being employed to combat this ancient scourge. By synthesizing key insights and highlighting gaps in knowledge, this review aims to inform policymakers, healthcare professionals, and researchers on the ongoing battle against "Malaria."

Keywords: Malaria Control, Disease Transmission, Public Health Strategies, Innovative Treatments

Introduction:

Malaria, a disease as ancient as civilization itself, continues to cast a long shadow over our modern world. Across continents and cultures, its insidious presence exacts a heavy toll on human lives, communities, and economies. In the intricate tapestry of global health, malaria stands as a formidable adversary, its impact reverberating far beyond the boundaries of affected regions (1).

The burden of malaria is staggering, with millions of cases reported annually and hundreds of thousands of lives lost, predominantly among the most vulnerable populations—children under five and pregnant women (2). Beyond the immediate human suffering, malaria exerts a profound socio-economic burden, impeding economic development, draining healthcare resources, and perpetuating cycles of poverty and inequality.

Amidst this backdrop of adversity, there exists an imperative for action—a collective resolve to confront malaria with all the tools at our disposal. This review seeks to shed light on the multifaceted nature of the malaria challenge, from its epidemiological complexities to the innovative strategies being employed in its control and elimination.

Objectives of the Review:

At its core, this review aims to serve as a compass, navigating the vast landscape of current research and control efforts in the realm of malaria. By synthesizing key findings from a diverse array of scientific literature, epidemiological studies, and field reports, our objective is to provide a comprehensive overview of the state of play in the fight against malaria.

Specifically, this review endeavors to achieve the following objectives:

Mapping the Epidemiological Landscape: We aim to delineate the global burden of malaria, highlighting regional variations in disease prevalence, transmission dynamics, and key

risk factors. By examining trends over time, we seek to identify areas of progress and persistent challenges in malaria control and elimination efforts (3).

Evaluating Control Strategies: From vector control interventions to antimalarial drug therapies, a multitude of strategies are employed in the battle against malaria. Our review will assess the efficacy, feasibility, and scalability of these interventions, drawing insights from both empirical research and real-world implementation experiences.

Identifying Knowledge Gaps and Research Priorities: Despite decades of research and concerted action, many aspects of malaria biology, transmission, and control remain poorly understood (4). Through a critical analysis of existing literature, we aim to identify key knowledge gaps and emerging research priorities that can inform future scientific inquiry and policy formulation.

Synthesizing Lessons Learned: Successes and setbacks in malaria control efforts offer valuable lessons for guiding future action. By synthesizing insights from diverse contexts and experiences, we seek to distill best practices, challenges, and opportunities for enhancing the effectiveness of malaria control and elimination strategies.

Epidemiology and Transmission Dynamics:

Malaria, caused by Plasmodium parasites and transmitted through the bites of infected Anopheles mosquitoes, continues to exact a heavy toll on human health and well-being worldwide. Understanding the epidemiology and transmission dynamics of malaria is crucial for designing effective control strategies and mitigating its impact on vulnerable populations (5).

Global Malaria Burden: Prevalence, Distribution, and Trends:

Malaria remains one of the most significant infectious diseases globally, with an estimated 229 million cases and 409,000 deaths reported in 2019 alone. While substantial progress has been made in reducing malaria incidence and mortality over the past two decades, the burden of the disease remains unevenly distributed, with sub-Saharan Africa bearing the greatest share of morbidity and mortality (5).

Within endemic regions, malaria transmission varies widely, influenced by factors such as altitude, climate, and human behavior. High transmission settings, characterized by intense year-round transmission, are prevalent in sub-Saharan Africa, while low transmission settings may exist in areas with seasonal or focal transmission patterns.

Despite significant gains in malaria control efforts, progress has plateaued in recent years, with several countries experiencing resurgences in malaria cases (6). Factors contributing to this stagnation include insecticide resistance, drug resistance, weakened health systems, and challenges in reaching remote and marginalized populations with essential malaria interventions.

Ecological Factors Influencing Malaria Transmission: From Vector Biology to Climate Change:

Malaria transmission is intricately linked to ecological factors that influence the abundance, behavior, and distribution of both mosquito vectors and human hosts. Anopheles mosquitoes, the primary vectors of malaria, exhibit diverse species-specific behaviors and preferences, with variations in biting habits, resting sites, and breeding habitats (7).

Vector control interventions, such as insecticide-treated bed nets and indoor residual spraying, target mosquito vectors at different stages of their life cycle, aiming to reduce their longevity, reproductive success, and biting rates. However, the efficacy of these interventions may

be compromised by factors such as insecticide resistance, environmental changes, and socio-economic disparities (8).



Climate change plays a significant role in shaping malaria transmission dynamics, influencing factors such as temperature, rainfall patterns, and ecological habitats conducive to mosquito breeding. Rising temperatures and altered precipitation patterns can extend the geographical range of malaria transmission, affect the seasonality of outbreaks, and create favorable conditions for mosquito proliferation (9).

Human Factors: Population Dynamics, Socioeconomic Disparities, and Migration Patterns:

Human population dynamics, socio-economic disparities, and migration patterns play critical roles in shaping malaria transmission and vulnerability. High population density, urbanization, and informal settlements can create conducive environments for malaria transmission, amplifying the risk of disease transmission in overcrowded and resource-limited settings (10).

Socioeconomic disparities, including poverty, limited access to healthcare, and inadequate housing, exacerbate the burden of malaria among marginalized and vulnerable populations. Lack of education, health literacy, and awareness about malaria prevention and control further contribute to disparities in disease burden and health outcomes.


Migration patterns, whether seasonal or permanent, can facilitate the spread of malaria across regions and continents, introducing new parasite strains and vectors to previously unaffected areas (11). Mobile and migrant populations, including refugees, displaced persons, and labor migrants, often face barriers to accessing healthcare and malaria prevention services, increasing their vulnerability to the disease.

Pathogenesis and Clinical Manifestations:

Understanding the pathogenesis and clinical manifestations of malaria is essential for accurate diagnosis, appropriate treatment, and effective management of the disease. Malaria, caused by Plasmodium parasites, manifests in a spectrum of clinical presentations, ranging from asymptomatic infection to severe and life-threatening complications (12).

The Biology of Plasmodium Parasites: Species Diversity and Virulence Factors:

Plasmodium parasites, belonging to the genus Plasmodium, comprise several species that are capable of infecting humans, including Plasmodium falciparum, Plasmodium vivax, Plasmodium malariae, Plasmodium ovale, and Plasmodium knowlesi. Each species exhibits distinct biological characteristics, including differences in morphology, lifecycle, and pathogenicity (13).



Among the Plasmodium species infecting humans, Plasmodium falciparum is the most virulent and responsible for the majority of malaria-related morbidity and mortality worldwide. Its ability to invade and multiply within red blood cells, as well as sequester in vital organs such as the brain and placenta, contributes to the severity of falciparum malaria infections (14).

Plasmodium vivax, while less virulent than P. falciparum, is characterized by its ability to form dormant liver stages (hypnozoites), leading to relapses months or even years after the initial infection. This unique feature poses challenges for malaria control and elimination efforts, particularly in regions where P. vivax is endemic (15).

Clinical Spectrum: From Asymptomatic Infection to Severe Malaria Complications:

Malaria presents along a spectrum of clinical manifestations, ranging from asymptomatic infection to uncomplicated malaria and severe malaria syndromes. Asymptomatic malaria

infections, characterized by the presence of Plasmodium parasites in the blood without overt symptoms, are common in endemic areas and contribute to the maintenance of transmission.

Uncomplicated malaria typically presents with symptoms such as fever, chills, headache, fatigue, and muscle aches. Other common symptoms may include nausea, vomiting, diarrhea, and jaundice. In uncomplicated cases, patients usually recover with appropriate antimalarial treatment and supportive care (16).

Severe malaria, on the other hand, represents a medical emergency and is associated with high morbidity and mortality rates, particularly in children under five and pregnant women. Severe malaria syndromes include cerebral malaria (characterized by altered consciousness and seizures), severe anemia, acute respiratory distress syndrome (ARDS), renal failure, and metabolic acidosis (17).

Host Factors Influencing Disease Outcomes: Genetic Susceptibility and Acquired Immunity:

Host factors play a crucial role in determining individual susceptibility to malaria infection and disease outcomes. Genetic factors, including polymorphisms in genes encoding red blood cell surface receptors (e.g., hemoglobin variants) and immune response genes (e.g., human leukocyte antigen alleles), influence an individual's susceptibility to malaria infection, severity of disease, and response to antimalarial treatment (18).

Acquired immunity, acquired through repeated exposure to malaria parasites over time, plays a central role in modulating disease severity and protecting against severe complications. Individuals living in endemic areas gradually develop immunity to malaria, characterized by a reduction in the frequency and severity of clinical episodes. However, this immunity is not

sterilizing and may wane in the absence of ongoing exposure, leading to an increased risk of severe disease upon re-infection (19).

Understanding the interplay between parasite biology, host factors, and clinical manifestations is essential for informing malaria diagnosis, treatment, and prevention strategies. By elucidating the mechanisms underlying malaria pathogenesis and immunity, researchers and healthcare providers can develop targeted interventions to reduce the burden of malaria and improve outcomes for affected populations (20).

Diagnosis and Surveillance:

Accurate diagnosis and effective surveillance are fundamental pillars of malaria control and elimination efforts. Rapid and reliable diagnostic tools enable early detection of malaria infections, facilitate appropriate treatment, and inform targeted interventions to prevent transmission and reduce morbidity and mortality. In this section, we explore the challenges in malaria diagnosis, recent advances in molecular diagnostics, and the role of surveillance systems in monitoring and controlling malaria (21).

Challenges in Malaria Diagnosis: Limitations of Microscopy and Rapid Diagnostic Tests

Microscopy, based on the examination of blood smears for the presence of Plasmodium parasites, has long been the gold standard for malaria diagnosis. While microscopy offers high specificity and the ability to quantify parasitemia, it has several limitations, including the requirement for skilled personnel, infrastructure, and time-consuming sample preparation and examination. Moreover, microscopy may miss low-density infections, leading to false-negative results, particularly in settings with limited resources and high transmission intensities (22).

Rapid diagnostic tests (RDTs), based on the detection of malaria antigens in blood samples, have revolutionized malaria diagnosis in resource-limited settings. RDTs are simple to use, provide rapid results within 15-20 minutes, and do not require specialized equipment or electricity. However, RDTs have limitations, including reduced sensitivity for detecting low-density infections, variability in performance between different brands and batches, and the inability to quantify parasitemia (23).

Advances in Molecular Diagnostics: PCR-Based Methods and Point-of-Care Technologies

PCR-based methods, including quantitative real-time PCR (qPCR) and loop-mediated isothermal amplification (LAMP), offer increased sensitivity and specificity for malaria diagnosis compared to conventional methods such as microscopy and RDTs. PCR-based assays can detect low-density infections, differentiate between *Plasmodium* species, and identify genetic markers associated with drug resistance and treatment failure (24).

Point-of-care (POC) molecular diagnostic technologies, such as nucleic acid amplification tests (NAATs) and portable PCR platforms, are emerging as promising tools for malaria diagnosis in remote and resource-limited settings. POC tests offer rapid results (within 1-2 hours), high sensitivity and specificity, and the potential for decentralized testing at the community level. Examples include the use of portable qPCR devices and cartridge-based LAMP assays for on-site detection of malaria infections (25).



Surveillance Systems: Monitoring Trends, Detecting Outbreaks, and Guiding Control Interventions

Surveillance systems play a critical role in monitoring malaria trends, detecting outbreaks, and guiding control interventions at local, national, and global levels. Traditional surveillance methods rely on passive case detection through health facility reporting and routine data collection systems. However, these systems may underreport cases, particularly in areas with weak health infrastructure and limited access to healthcare (26).

Enhanced surveillance approaches, including active case detection, sentinel surveillance, and mobile health (mHealth) technologies, complement traditional methods and improve the timeliness, completeness, and accuracy of malaria data. Real-time reporting systems, geographic

information systems (GIS), and predictive modeling techniques enable early detection of outbreaks, targeted interventions, and allocation of resources to high-risk areas (27).

Integrated surveillance platforms, such as the World Health Organization's (WHO) Global Malaria Programme and national malaria control programs, facilitate data sharing, coordination, and collaboration between stakeholders involved in malaria control and elimination efforts (28). These platforms provide a framework for monitoring progress towards malaria targets, evaluating the impact of interventions, and adapting strategies in response to changing epidemiological patterns and emerging challenges.

Malaria prevention and control require a multi-faceted approach that addresses both the human and environmental factors contributing to transmission. In this section, we explore key strategies for malaria prevention and control, including vector control interventions, chemoprophylaxis and treatment guidelines, and integrated approaches involving community engagement and multi-sectoral collaboration (29).

Vector Control Interventions: Insecticide-Treated Bed Nets, Indoor Residual Spraying, and Larval Control

Vector control interventions aim to reduce mosquito populations and minimize human-vector contact, thereby interrupting malaria transmission. Insecticide-treated bed nets (ITNs) and long-lasting insecticidal nets (LLINs) are widely used as frontline tools for malaria prevention, providing physical barriers against mosquito bites and delivering insecticidal effects to kill mosquitoes (30).

Indoor residual spraying (IRS) involves the application of insecticides to the interior walls of houses and other indoor resting surfaces to kill mosquitoes that come into contact with treated

surfaces. IRS is particularly effective in areas where Anopheles mosquitoes prefer to rest indoors after feeding on blood.

Larval control measures target mosquito breeding sites, such as stagnant water bodies and open drains, to reduce mosquito populations before they emerge as adults. Larviciding, habitat modification, and environmental management strategies aim to disrupt mosquito breeding habitats and limit larval development.

Chemoprophylaxis and Treatment Guidelines: Evolving Recommendations and Drug Resistance Concerns

Chemoprophylaxis involves the use of antimalarial drugs to prevent malaria infection in individuals at risk of exposure, such as travelers to malaria-endemic regions and pregnant women living in areas of high transmission. Recommended drugs for chemoprophylaxis vary depending on factors such as the destination, Plasmodium species prevalence, and drug resistance patterns.

Treatment guidelines for malaria encompass the selection of appropriate antimalarial drugs based on factors such as the species of Plasmodium parasite, disease severity, patient age, pregnancy status, and drug resistance profiles. Artemisinin-based combination therapies (ACTs) are the first-line treatment for uncomplicated falciparum malaria, while other antimalarial drugs may be used for non-falciparum species or severe malaria cases (31).

Concerns about drug resistance, particularly to artemisinin derivatives, pose significant challenges to malaria control and treatment efforts. Surveillance for antimalarial drug resistance, including molecular monitoring of resistance markers and therapeutic efficacy studies, is essential for detecting emerging resistance patterns and informing treatment policies.

Integrated Approaches: Community Engagement, Health Education, and Multi-Sectoral Collaboration

Integrated approaches to malaria prevention and control involve engaging communities, empowering individuals, and fostering collaboration across sectors to address the social, economic, and environmental determinants of malaria transmission. Community engagement strategies promote ownership, participation, and sustainability of malaria control interventions, involving community leaders, health workers, and local stakeholders in decision-making processes (32).

Health education and behavior change communication initiatives raise awareness about malaria prevention methods, symptoms, and treatment-seeking behaviors, empowering individuals to take proactive steps to protect themselves and their families from malaria. Information, education, and communication (IEC) materials, interpersonal communication, and mass media campaigns are used to disseminate key messages and promote behavior change.

Multi-sectoral collaboration brings together diverse stakeholders, including government agencies, non-governmental organizations (NGOs), academic institutions, and private sector partners, to leverage resources, expertise, and networks in support of malaria control efforts. Coordinated action across sectors such as health, education, agriculture, water and sanitation, and environmental management enhances the impact and sustainability of malaria interventions (33).

Conclusion:

Malaria remains a formidable global health challenge, with millions of cases reported annually and significant morbidity and mortality, particularly among vulnerable populations in low-resource settings. Despite the complex and dynamic nature of malaria transmission,

significant progress has been made in recent decades through concerted efforts in prevention, diagnosis, and treatment.

Vector control interventions, including insecticide-treated bed nets, indoor residual spraying, and larval control, have played a crucial role in reducing malaria transmission by targeting mosquito vectors and minimizing human-vector contact. These interventions, when implemented effectively and in combination, have proven to be highly effective in reducing malaria incidence and mortality, particularly in endemic regions.

However, the emergence of drug resistance and insecticide resistance poses significant challenges to malaria control efforts, threatening the efficacy of current prevention and treatment strategies. Continued surveillance for drug resistance and vector resistance, along with the development and deployment of alternative interventions, are critical to maintaining progress and adapting to evolving malaria dynamics.

Chemoprophylaxis and treatment guidelines provide essential tools for preventing malaria infection in high-risk populations and ensuring prompt and effective treatment for individuals with malaria. Ongoing research into new antimalarial drugs, vaccines, and diagnostic technologies offers hope for improved tools and strategies for malaria control and elimination in the future.

Integrated approaches, grounded in community engagement, health education, and multi-sectoral collaboration, are essential for addressing the socio-economic determinants of malaria transmission and promoting sustainable solutions. By empowering communities, fostering collaboration across sectors, and leveraging resources and expertise, we can strengthen malaria control efforts and accelerate progress towards a malaria-free world.

The fight against malaria requires a holistic and multi-faceted approach, encompassing prevention, diagnosis, treatment, and community engagement. With continued investment, innovation, and collaboration, we have the opportunity to overcome the challenges posed by malaria and realize the vision of a world where malaria is no longer a threat to human health and well-being.

Reference:

1. World Health Organization . *World Malaria Report 2018*. WHO; Geneva, Switzerland: 2018.
2. White N.J.N., Pukrittayakamee S., Hien T.T.T., Faiz M.A., Mokuolu O.A.O., Dondorp A.A.M. Malaria. *Lancet*. 2014;383:723–735. doi: 10.1016/S0140-6736(13)60024-0.
3. Pan American Health Organization. World Health Organization . *Epidemiological Alert, Increase of Malaria in the Americas*. PAHO; Washington, DC, USA: 2018.
4. Dhiman S. Are malaria elimination efforts on right track? An analysis of gains achieved and challenges ahead. *Infect. Dis. Poverty*. 2019;8:14. doi: 10.1186/s40249-019-0524-x.
5. Walker N., Nadjm B., Whitty C. Malaria. *Medicine*. 2017;42:52–58. doi: 10.1016/j.mpmed.2013.11.011.
6. Antinori S., Galimberti L., Milazzo L., Corbellino M. Biology of human malaria plasmodia including Plasmodium knowlesi. *Mediterr. J. Hematol. Infect. Dis*. 2012;4:e2012013. doi: 10.4084/mjhid.2012.013.
7. Ashley E.A., Pyae Phyo A., Woodrow C.J. Malaria. *Lancet*. 2018;391:1608–1621. doi: 10.1016/S0140-6736(18)30324-6.
8. Singh B., Daneshvar C. Human Infections and Detection of Plasmodium knowlesi. *Clin. Microbiol. Rev*. 2013;26:165–184. doi: 10.1128/CMR.00079-12.

9. Vuk I., Rajic Z., Zorc B. Malaria and antimalarial drugs. *Farm Glas.* 2008;64:51–60.
10. Soulard V., Bosson-Vanga H., Lorthiois A., Roucher C., Franetich J.-F., Zanghi G., Bordessoulles M., Tefit M., Thellier M., Morosan S., et al. Plasmodium falciparum full life cycle and Plasmodium ovale liver stages in humanized mice. *Nat. Commun.* 2015;6:7690. doi: 10.1038/ncomms8690.
11. Josling G.A., Llinás M. Sexual development in Plasmodium parasites: Knowing when it's time to commit. *Nat. Rev. Genet.* 2015;13:573–587. doi: 10.1038/nrmicro3519.
12. Blut A. Untergruppe "Bewertung Blutassoziierter Krankheitserreger." Malaria. *Transfus. Med. Hemother.* 2009;36:48–60. doi: 10.1159/000197327.
13. Cowman A., Healer J., Marapana D., Marsh K. Malaria, Biology and Disease. *Cell.* 2016;167:610–624. doi: 10.1016/j.cell.2016.07.055.
14. Cartwright F., Biddis M. *Disease and History*. Naklada Ljevak; Zagreb, Croatia: 2006.
15. Lehrer S. *Vitezi Medicine*. Prosveta; Beograd, Serbia: 1981.
16. Moss W., Shah S., Morrow R. The History of Malaria and its Control. *Int. Encycl. Public Health.* 2008:389–398. doi: 10.1016/B978-012373960-5.00374-9.
17. Dugacki V. Dr. Rudolf Battara operation in Nin in 1902, the first systematic battle attempt against malaria in Croatia. *Med. Jaderina.* 2005;35:33–40.
18. Jovic S. *Istorija Medicine i Zdravstvene Kulture na tlu Dansnje Vojvodine 1718–1849 II. dio*. Matica srpska, Srpska akademija nauka i umetnosti; Novi Sad, Serbia: 1998.
19. Tan S.Y., Ahana A. Charles Laveran (1845–1922), Nobel laureate pioneer of malaria. *Singap. Med. J.* 2009;50:657–658.
20. Saifi M.A., Siddiqui M.I., Al-Khalifa M.S. Malaria: Patterns of relapse and resistance. *J. King Saud Univ. Sci.* 2010;22:31–36. doi: 10.1016/j.jksus.2009.12.005.

21. Chu C.S., White N.J. Management of relapsing Plasmodium vivax malaria. *Expert Rev. Anti-Infect. Ther.* 2016;14:885. doi: 10.1080/14787210.2016.1220304.
22. Murphy S.C., Shott J.P., Parikh S., Etter P., Prescott W.R., Stewart V.A. Malaria diagnostics in clinical trials. *Am. J. Trop. Med. Hyg.* 2013;89:824–839. doi: 10.4269/ajtmh.12-0675.
23. Tangpukdee N., Duangdee C., Wilairatana P., Krudsood S. Malaria diagnosis, a brief review. *Korean J. Parasitol.* 2009;47:93–102. doi: 10.3347/kjp.2009.47.2.93.
24. Trampuz A., Jereb M., Muzlovic I., Prabhu R.M. Clinical review, Severe malaria. *Crit. Care.* 2003;7:315–323. doi: 10.1186/cc2183.
25. She R.C., Rawlins M.L., Mohl R., Perkins S.L., Hill H.R., Litwin C.M. Comparison of Immunofluorescence Antibody Testing and Two Enzyme Immunoassays in the Serologic Diagnosis of Malaria. *J. Travel Med.* 2007;14:105–111. doi: 10.1111/j.1708-8305.2006.00087.x.
26. Oh J.S., Kim J.S., Lee C.H., Nam D.H., Kim S.H., Park D.W., Lee C.K., Lim C.S., Park G.H. Evaluation of a malaria antibody enzyme immunoassay for use in blood screening. *Mem. Inst. Oswaldo Cruz.* 2008;103:75–78. doi: 10.1590/S0074-02762008005000008.
27. Kakkilaya B. Rapid Diagnosis of Malaria. *Lab. Med.* 2003;34:602–608. doi: 10.1309/J4ANKCCJ147JB2FR.
28. Abba K., Kirkham A.J., Olliaro P.L., Deeks J.J., Donegan S., Garner P., Takwoingi Y. Rapid diagnostic tests for diagnosing uncomplicated non-falciparum or Plasmodium vivax malaria in endemic countries. *Cochrane Database Syst. Rev.* 2014;12 doi: 10.1002/14651858.CD011431.

29. World Health Organization . *Guidelines for the Treatment of Malaria*. 3rd ed. WHO; Geneva, Switzerland: 2015.
30. World Health Organization . *Malaria Rapid Diagnostic Test Performance, Results of WHO Product Testing of Malaria RDTs, Round 8 (2016–2018)* WHO; Geneva, Switzerland: 2018.
31. Ranadive N., Kunene S., Darteh S., Ntshalintshali N., Nhlabathi N., Dlamini N., Chitundu S., Saini M., Murphy M., Soble A. Limitations of Rapid Diagnostic Testing in Patients with Suspected Malaria, A Diagnostic Accuracy Evaluation from Swaziland, a Low-Endemicity Country Aiming for Malaria Elimination. *Clin. Infect. Dis.* 2017;64:1221. doi: 10.1093/cid/cix131.
32. Berhane A., Russom M., Bahta I., Hagos F., Ghirmai M., Uqubay S. Rapid diagnostic tests failing to detect *Plasmodium falciparum* infections in Eritrea, an investigation of reported false negative RDT results. *Malar. J.* 2017;16:105. doi: 10.1186/s12936-017-1752-9.
33. Kozycki C.T., Umulisa N., Rulisa S., Mwikarago E.I., Musabyimana J.P., Habimana J.P., Karema C., Krogstad D.J. False-negative malaria rapid diagnostic tests in Rwanda: Impact of *Plasmodium falciparum* isolates lacking *hrp2* and declining malaria transmission. *Malar. J.* 2017;16:123. doi: 10.1186/s12936-017-1768-1.